

Tel: +420 603 476 934 **E-mail:** top-bio@top-bio.cz

www.top-bio.com

CERTIFICATE OF ANALYSIS

Product: Carrier-ACRYL

Catalog No: C081, C082

Lot No: C081122026

Date of Expiry: 12/2026

Composition: Linear polyacrylamide in deionized and sterile water (18 Mohm.cm) at concentration ~25

mg/ml.

Storage temperature: For short terms (months) at temperature +4°C ± 3°C. For long terms at temperature -20 ±

5°C. Material can be repeatedly defrosted.

UV (Abs. 10 mm) 220 nm = 21 ± 2; 240 nm = <0.1; 260 nm < 0.1.

Functional Test: Each batch of Carrier-ACRYL is analyzed in several assays. For the assays, DNA or RNA is

examined in the Carrier Assay Buffer (CAB): 10 mM Tris-HCl, 2 mM MgCl2, 1 mM

dithiothreitol, pH 7.5 at 37°C.

Nucleic acid precipitation assay. Economy DNA marker (Cat. No. D071, $2.5~\mu$ l) is mixed with 0.2 ml 10 mM Tris buffer, pH 8.0 + 1 mM EDTA, 1 μ l Carrier-ACRYL, 20 μ l of 3 M sodium acetate, pH 5.2, and 0.6 ml of 96% Ethanol. After 30 minutes at 2 - 8°C the mixture is centrifuged for 10 min at 12,000 x g, analyzed by electrophoresis in agarose gel with ethidium bromide and observed under UV light. More than 90% of all components of the DNA marker is recovered in the precipitate.

Nick activity assay. Plasmid pUC19 (1 μ g) in 50 μ l CAB with Carrier-ACRYL (50 μ g) is incubated for for 1 hour at 37°C, followed by electrophoresis in agarose gel with ethidium

bromide. No nicking activity is observed.

Ribonuclease assay. RNA (1 μ g) in 50 μ l CAB with Carrier-ACRYL (50 μ g) is incubated for 1 hour at 37°C, followed by electrophoresis in agarose gel with ethidium bromide. No

changes in properties of RNA are observed under UV light.

Result: passed

FOR RESEARCH USE APPROVED DATE: 13. 12. 2024

Manager: Hana Těšitelová